

Okayama, January 29, 2019

Exercises for Advanced Physics 3, 2018 term 4

Exercise Set 6

(Due date: Tuesday, February 5, 2019)

Exercise 6 (Interchange of spins) (10 points)

We consider the Pauli spin operator $\vec{\sigma} = (\sigma_x, \sigma_y, \sigma_z)$ with

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

that obey the relations

$$(1) \quad \begin{aligned} \sigma_i^2 &= 1, \quad i = x, y, z \\ \sigma_i \sigma_j &= i \sigma_k, \quad (i, j, k) \in \{(x, y, z), (z, x, y), (y, z, x)\} \\ [\sigma_i, \sigma_j]_+ &= \sigma_i \sigma_j + \sigma_j \sigma_i = 2 \delta_{ij} \mathbb{1}, \quad \mathbb{1} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \end{aligned}$$

$\vec{\sigma}^{(i)}$, $i = 1, 2$ is the spin operator for the particle i .

(a) Prove $(\vec{\sigma}^{(1)} \cdot \vec{\sigma}^{(2)})^2 = 3\mathbb{1} - 2(\vec{\sigma}^{(1)} \cdot \vec{\sigma}^{(2)})$

(b) Show that the operator

$$Q_{12} = \frac{1}{2}(\mathbb{1} + \vec{\sigma}^{(1)} \cdot \vec{\sigma}^{(2)})$$

interchanges the spins of particles 1 and 2:

$$(2) \quad \begin{aligned} Q_{12} \vec{\sigma}^{(1)} Q_{12}^{-1} &= \vec{\sigma}^{(2)} \\ Q_{12} \vec{\sigma}^{(2)} Q_{12}^{-1} &= \vec{\sigma}^{(1)} \end{aligned}$$