Okayama University Faculty of Science

Research Institute for Interdisciplinary Science Prof. Harald Jeschke Tutor: Makoto Shimizu



Okayama, December 10, 2019

Exercises for Advanced Physics 2, 2019 term 4

Exercise Set 1 (Due date: Tuesday, December 17, 2019)

Exercise 1 (Landau diamagnetism) (10 points)

Consider the two-dimensional electron gas in the presence of a perpendicular field $\vec{B}_0 = B_0 \vec{e}_z$. In the plane, assume that the electron gas is enclosed in a rectangular sample with side lengths L_x and L_y . According to section 3.3.1 of the script, in the ground state, the N electrons occupy the Landau levels

(1)
$$E_{n,k_x} = \hbar \omega_c \left(n + \frac{1}{2}\right), \quad n = 0, 1, 2, \dots$$
$$\omega_c = \frac{eB_0}{m}$$

The spin splitting is neglected here.

- (a) What is the smallest field $B_0 = B_0^{(0)}$ at which all the electrons are placed in the n = 0 level?
- (b) What is the field $B_0 = B_0^{(n_0)} \leq B_0^{(0)}$ at which the N electrons are uniformly distributed in the Landau levels up to the quantum number n_0 ?
- (c) If the field B_0 is between the two critical fields $B_0^{(n_0)}$ and $B_0^{(n_0-1)}$

$$\mathsf{B}_0^{(\mathfrak{n}_0-1)} \geqslant \mathsf{B}_0 \geqslant \mathsf{B}_0^{(\mathfrak{n}_0)},$$

calculate the total energy $E(B_0)$ of the N electron system.

(d) What is the result for the special case $E(B_0^{(n_0)})$?

Please explain all steps!